



The 6<sup>th</sup> international- 20<sup>th</sup> Arabic conference for  
Home Economics  
Home Economics and Educational quality assurance  
December 23rd -24th, 2018

<http://homeEcon.menofia.edu.eg>

ISSN 1110-2578

**Journal of Home  
Economics**

## **Protective effect of banana and mango peels against lead toxicity in rats**

**\*Ereny Wilson Nagib, \*Hala RashedAtaya**

\*Department of Home Economic, Faculty of Specific Education, Ain Shams University,  
Cairo, Egypt.

### **Abstract**

This study was carried out to determine the effect of dry banana peels (*Musa Acuminata*) and drymango peels (*Mangifera indica L.*) against lead toxicity in weaning rats. Forty five growing male albino were divided into two main groups. The first group was the negative group (9) rats, the second group (36) rats was exposed to lead acetate and divided into four groups. The subgroup one positive group fed basal diet, the subgroup two fed basal diet containing (15%) banana peels, the subgroup three fed basal diet containing (15%) mango peels, the subgroup four fed basal diet containing (15%) combination of (banana and mango peels) for (6) weeks.

At the end of the experimental period the best result found in group fed basal diet containing (15%) combination banana and mango peels improved daily feed intake, body weight gain, feed efficiency ratio, hemoglobin and serum iron level. Moreover, lead concentration was decreased in serum and also it was improved the concentration of calcium, phosphorous as well as lead in bone of intoxicated rats. Dry peels of fruits were improved liver enzymes and kidney functions compared with positive group.

The histopathology results were agreed with the results of serum parameters. These results recommend increasing the consumption of banana and mango peels. Nutrition Education Programs are needed to illustrate the importance of banana and mango peels in reducing lead toxicity.

**Key words:** lead toxicity, banana, mango peels.

## **Introduction**

Lead (Pb) is one of the pervasive and persistent environmental toxic metals. Despite considerable efforts to identify and eliminate sources of pbexposure, this metal remains induces adverse health effects for centuries (**khodamoradi et al., 2015**). According to **World Health Organization (WHO) statisticsthe Institute for Health Metrics and Evaluation (IHME) has estimated that in (2013)**accounted lead exposure for 853 000 death due to long-term effects on health, with the highest burden in low and middle income countries. IHME also estimated that lead exposure accounted for 9.3% of the global burden of idiopathic intellectual disability, 4% of the global burden of ischemic heart disease 6.6% of the global burden of stroke. (Pb)had modest uses in ancient medicines. Today, it used in many products including alkyl-lead petroleum combustion, production, and storage of lead-acid batteries, leaded glass, cement manufacture, production of plastics & ceramics and also found in some imported cosmetics (**karamian et al., 2015**). Lead (Pb)can damage various body systems; however, the central nervous system is the primary target (**karamian et al., 2015 and Li et al., 2016**). Several reports have demonstrated that exposure to lead causes deleterious effects on the nervous system, including decrements in iq, impaired cognition, and memory, as well as impaired peripheral nerve functions and related behavioral disturbances (**khodamoradi et al., 2015**). The developmental toxicity of lead has become a significant area of research since children are much more sensitive than adults to learning impairment following low level of lead exposure (**Davis and Svendsgaard, 1987**).

Lead (Pb) poisoning negatively influenced various body systems particularly, hematopoietic and renal system (**Ercal et al., 1996**). Gastrointestinal, reproductive, circulatory, and immunological pathologies (**patrick, 2006**), reproductive dysfunctions, (**Marchlewicz et al., 1993**). Moreover, lead inhibition of the activities of antioxidant enzymes, including glutathione peroxidase, catalase and superoxide dismutase (**Silbergeld et al., 2000**). Furthermore, stimulation of lipid peroxidation and depletion of antioxidant reserves which was postulated to be major contributors to lead-exposure related diseases (**Patrick, 2006**).

Banana(*Musa Acuminata*) The peel could be a potential source of antioxidant and antimicrobial activities. Ethyl acetate and water soluble fractions of green banana peel displayed high antimicrobial and antioxidant activities. Most of the compounds isolated from green peel,  $\beta$ -sitosterol, malic acid, 12-hydroxystearic acid and succinic acid, showed significant antibacterial activities and low antioxidant activities, while those compounds isolated from water soluble extracts, glycoside and monosaccharide components, displayed significant antioxidant and low antimicrobial activities (**Matook and Fumio, 2005**). Banana peel represents about 40% of total weight of the fresh fruit (**Anhwange et al., 2008**). The total amount of phenolic compounds in banana peel has been ranged from 0.90 to 3.0 g/100 g dry weight and gallic acid is identified at a concentration of 160 mg/100 g dry weight; **Someya et al., (2002)**. Other phytochemicals such as anthocyanin, delphinidin, cyaniding; Seymour, and catecholamines have been identified (**Kanazawa and Sakakibara, 2000**) in ripe banana pulp and peel. Recent studies demonstrated that banana peel generally includes higher phenolic compounds than those of banana pulps; **Kondo et al. (2005) and Sulaiman et al. (2011)**. **Subagio et al. (1996)** identified carotenoids such as  $\beta$ -carotene,  $\alpha$ -carotene and different xanthophylls in the range of 300– 400  $\mu$ g/100 g of banana peels. **Gonzalez- Montelongo et al. (2010)** studied the extraction conditions that produce maximum antioxidant activity (Acetone: water (1:1), 25 °C, 120 min). Moreover, the number of extraction steps, temperature and time, have been reported as the most effective factors associated with antioxidant properties of banana peel, respectively.

Mango(*Mangifera indica L.*) is one of the most important tropical fruits worldwide in terms of production and consumer acceptance (**FAO, 2012**). Mango byproducts contain significant amounts of phytochemicals, which makes them suitable to be processed for value-added applications in functional foods and nutraceuticals (**Ajila et al., 2007, 2010**). Mango peel contains various classes of polyphenols, carotenoids, and vitamins with different health-promoting properties, mainly antioxidant activity (**Schieber et al., 2003; Ajila et al., 2008; Manthey & Perkins-Veazie, 2009**).

Therefore, the present work was conducted to study the protective effect of banana and mango peels against lead toxicity in rats.

### **Materials and Methods**

#### **Materials:**

Banana (*Musa Acuminata*) and mango (*Mangifera indica* L.) were purchased from local market in Egypt.

Albino rats (Sprague - Dawley Strain) weighting ( $70 \pm 5$ g) were purchased from Helwan Experimental Animals Station.

Casein, vitamins, minerals, cellulose, choline chloride obtained and pure lead acetate were purchased from El-Gomhorya Company, Cairo, Egypt.

Kits measurement of all parameters for biological experimental were purchased from BiconDiagnosemittel GmbH and Co. KG Hecke 8 made in Germany.

#### **Methods:**

##### **Banana and mango peels preparation**

The peels were removed from the fresh and submitted to the following processes:

1. Fresh peels: banana and mango fresh peels were washed with water and cut into small pieces.
2. Dried peels: peel samples were dried at 50 °C in a hot air oven for 12 h and ground to obtain the particle size of less Than 1.0 mm (**Adejuyitan et al., 2008**).

##### **Chemical composition analysis:**

Moisture, total protein, total lipid and fiber were analyzed in banana and mango peels according to **AOAC (2010)**.

##### **Determination of total phenolic compounds:**

The samples were extracted in terms of the method described by **Garcia-Salas et al. (2010)** and used for spectrophotometric analyses. The total phenolic content of the extracts was determined by a Folin-Ciocalteu phenol reagent method (**Xu and Chang, 2007**) using gallic acid as standard. The absorbance was measured with UV/vis spectrophotometer (Varian Cary 50 Scan, Australia) at 760 nm. A mixture of 80% methanol and reagents was used as a blank solution. The total phenolic content was expressed as

gallic acid equivalents ( $\mu\text{g}$  of GAE/mg dry matter) through the calibration curve of gallic acid.

#### **Mineral composition of fruit peels**

The mineral composition of the banana and mango peels was determined according to the methods of the **A.O.A.C. (2005)**.

#### **Biological investigations:**

Albino rats forty five (Sprague – Dawley strain) of weaning rats, weighting  $70 \pm 5$  g each were housed in an individual stainless steel cage under hygienic controlled condition. The basal diet was prepared according to (**Reeves et al., 1993**) after this period, the rats were divided into two main groups, as follows: The first main group (9 rats) was fed on the basal diet for another six weeks (42 days) and considered as negative control. The second main group (36 rats) were exposed to lead acetate at (200 mg/ kg from weight rat) daily drinking orally according to (**Newairy and Abdou., 2009**) and divided into four sub groups (9 rats for each). The first one (9 rats) was continued to be fed on basal diet and considered as positive control. From the second to four subgroups (9 rats for each) were fed on basal diet fortified with 15% separately from dried banana, mango peels and mixed from them, respectively. During the experimental period (6 weeks), the diets consumed and body weights were recorded twice weekly and feed efficiency ratio according to (**Chapman et al., 1959**).

At the end of the experiment, the animals were fasting overnight, then the rats were anaesthetized and sacrificed, and blood samples were collected. Blood samples were centrifuged and the serum was separated to estimate some biochemical parameters. Blood haemoglobin (Hg) was determined using enzymatic calorimetric method was described by **Young, (1990)**. Lead and iron concentration in serum was determined according to **Parson (2001) and Ramsay (1957)**. Tibia and organs as liver, kidney were weighted. Moreover, calcium, phosphorus and lead in bone were determined according to **Carter and Gregorich (2006)**. The samples were digested using a mixture of nitric and perchloric acid at ratio of 3:1, then P, Ca and Pb were determined by using atomic absorption spectrophotometer. Liver functions (AST and ALT enzymes) was estimated by **Sherwin (1984)** and

kidney function as uric acid, urea and creatinine were determined according to **Haisman and Muller (1977)**, **Henry et al. (1974)** and **Larsen (1972)**,  
**Statistically analysis:**

The data obtained in the present study was analyzed by ANOVA. For all analyses, when a significant difference ( $p < 0.05$ ) was detected in some variable, the data means test was applied to evaluate the difference between the samples. The results were analyzed with the aid of the software SAS System for Windows **SAS (2008)**.

### **Results and Discussion**

#### **Chemical Composition of banana and mango peels:**

The results presented in Table (1) shows that the carbohydrate, fiber and protein content of dried banana, mango peels were the highest 63.0, 13.70 and 6.25% in dried banana peels, meanwhile; mango peels were 61.70, 17.33 and 3.61%, respectively. Whereas, the total lipids and ash content were recorded 7.70 and 2.50 in dried banana and also it was 4.23 and 4.88% in dried mango peels, respectively. **HappiEmaga et al. (2007)** showed that the banana peel had higher fat, ash, and total dietary fiber content, but lower protein and starch content than those of the banana peels **Figuerola et al. (2005)**, **Llobera and Canellas (2007)** and **Marin et al. (2007)** reported by had lower total dietary fiber content than fiber obtained from different sources of fruit industrial by-products (60–78 g/100 g dry matter).

**Ashoush and Gadallah (2011)** studied the effect of mango peels powders (MPP) at different replacing levels (5, 10, 15 and 20%) and mango kernels powders (MKP) at (20, 30, 40 and 50%) on rheological, physical, sensory and antioxidant properties of biscuits were evaluated. The results revealed that MPP had high contents of ash, crude fiber and water holding capacity

(**Serafini et al., 1998** and **Carbonnea et al., 1998**). Consumption of controlled diets high in fruits and vegetables increased significantly the antioxidant capacity of plasma, and the increase could not be explained by the increase in the plasma R-tocopherol or carotenoid concentration (**Cao et al., 1998**). Moreover, epidemiological studies have found that there is a significant negative association between the intake of fruits and vegetables and heart disease mortality (**Hertog et al., 1993, 1995; Knekt et al., 1996**).

Flavonoids and other plant phenolics, such as phenolic acids, stilbenes, tannins, lignans, and lignin, are especially common in leaves, flowering tissues, and woody parts such as stems and barks (**Larson, 1988**). The antioxidant activity of phenolics is mainly due to their redox properties, which allow them to act as reducing agents, hydrogen donors, and singlet oxygen quenchers. In addition, they have a metal chelation potential (**Rice-Evans et al., 1995**).

According to the study by **Someya et al. (2002)** total phenolics are more abundant in peel (907 mg/ 100 g dry wt.) than in pulp (232 mg/100 g dry wt.)

**Sogi et al., (2013)** evaluate different drying methods for mango peel and kernel and assess their impact on nutrients, antioxidants and functional properties. Mango peel and kernel were dried using different techniques, such as freeze drying, hot air. Freeze dried mango waste had higher antioxidant properties than those from other techniques. The total phenolics content of dried peel and kernel powders were 2032–3185 and 11,228–20,034 mg GAE/100 g respectively. The cabinet dehydrated mango waste can be utilized in several food applications due to their content of phytochemicals that exert antioxidant properties.

**Table (1): Chemical composition of dry banana and mango peels (g / 100g):**

<b>Fruits nutrients</b>	<b>Banana peels</b>	<b>Mango peels</b>
Moisture	6.70 ± 02.22	4.92 ± 0.32
Protein	6.25± 0.25	3.61 ± 0.15
Total lipid	7.70 ± 0.10	4.23 ± 0.10
Total carbohydrates	63.00 ± 1.36	61.70 ± 0.15
Fiber	13.70 ± 0.25	17.33 ± 0.61
Ash	2.50 ±1.52	4.88 ± 0.59
Total phenolic (µg of GAE/g)	10.90± 0.10	23.06 ± 0.40

The data were presented as mean±S.D.

**Mineral composition of dry banana and mango peels (g / 100g):**

Minerals play a key role in various physiological functions of the body, especially in the building and regulation processes. Fruits are good source of

dietary minerals (Ismail *et al.*,2011). The mineral composition of fruit peels is represented in table (2) show that mango peels are high content of calcium 60.63mg and banana peels are high in iron 15.15 mg.

Adlin (2008) and Faigin (2001) reported that the content of a banana peel was 55.59%, calcium 0.36%, Phosphor 0.10% and gross energy 3727 kcal/kg. Banana peel also contains vitamins C, E, and B6. Vitamin C can act as an antioxidant, while serotonin is thought to play an anti-depressant so as to increase feed intake and body weight on heat stress conditions.

**Table (2): Mineral composition of dry banana and mango peels (mg / 100g):**

Elements (mg/100g dry peel)	Fruits	
	banana	Mango
<b>calcium</b>	19.86 ± 0.24	60.63± 4.58
<b>zinc</b>	1.72 ± 0.17	0.66 ± 0.06
<b>Iron</b>	15.15 ± 0.36	12.79±1.56
<b>Manganese</b>	9.05 ± 0.34	4.77 ± 0.22

Values are means ± standard deviations of three independent analyses.

**Body weight gain, total feed intake and feed efficiency ratio of rats suffering from lead toxicity.**

The results in Table (3) showed that the control positive reduction in feed intake, body weight gain and feed efficiency when rats exposed to lead toxicity (-8.0, 5.48 and -0.44) compared to negative control group was 13.01, 7.96 and 0.62, respectively. While rats were fed separately 15 % banana peels, mango peels and mixture banana and mango peels fortified with basal diet showed that significant increasing values compared with the positive group and the formula 3 (15% mixture with banana and mango peels) give the best result. The present results are supported by the study of the Mingeum Jeong *et al.* (2015) who reported that body weight of rats treated with 17 $\alpha$ -ethinyloestradiol for 6 weeks were lower than that measured in the control group. Dietary guidelines have recommended increased intakes of



fruits and vegetables (Willett et al., 1995). The rationale for such a hypothesis of weight loss with increased intake of fruits is based on three premises: the low-energy density of most fruits, their higher fiber composition and a less striking variation of diets high in fruit. In support of this hypothesis, (Howarth et al., 2001) indicated that, under fixed energy intake, soluble or insoluble fiber intake increases post meal satiety and decreases subsequent hunger. In addition, this review suggested, at least for short-term follow-up, that high fiber diets decrease energy intake and body weight. Fruit consumption has a potential role in the prevention of overweight and obesity ( Tetens and Alinia, 2009).

**Table (3): Body weightgain,total feed intakeand feed efficiency ratio of rats suffering from lead toxicity.**

Groups	Initial body weight (g)	Body weight gain (g)	Total feed intake (g/day)	Feed efficiency ratio
Control negative	74.0 ± 2.60 <sup>a</sup>	15.01±2.25 <sup>a,b</sup>	7.96±1.02 <sup>b</sup>	0.53±1.02 <sup>a</sup>
Control positive	71.2 ± 3.58 <sup>a</sup>	- 8.0±1.30 <sup>d</sup>	5.48±1.01 <sup>c</sup>	-0.44±0.03 <sup>b</sup>
Group 1	73.2±2.44 <sup>a</sup>	12.93±2.21 <sup>a,b</sup>	7.18±0.97 <sup>b</sup>	0.45±0.07 <sup>a,b</sup>
Group 2	72.8 ± 3.47 <sup>a</sup>	13.33±0.77 <sup>a,b</sup>	7.28±0.87 <sup>b</sup>	0.47±0.06 <sup>a,b</sup>
Group 3	69.4±9.85 <sup>a</sup>	19.31±2.97 <sup>a</sup>	9.91±1.01 <sup>a</sup>	0.48±0.92 <sup>a,b</sup>

Values are expressed as mean ± SD.- Significant at p<0.05 using one way ANOVA test.- Values which have different letters in each column differ significantly, while those with have similar or letters are not significant

**Weight of relative organs rats suffering from lead toxicity**

Table (4) revealed that the results of mean weight value the organs as liver, kidney and tibia. These organs were increased when rats exposed to lead toxicity (control positive) 5.12, 1.21 and 1.93 %, respectively compared with negative control group was 2.92, 1.08 and 0.89%, respectively.

**Table (4): Percent of relative organs weight in rats suffering from lead toxicity.**

.Groups	Weigh of organs rats		
	Liver (%)	Kidney (%)	Tibia (%)
Control negative	2.92±0.42 <sup>b</sup>	1.08±0.24 <sup>a</sup>	0.89±0.08 <sup>b</sup>
Control positive	5.12±0.22 <sup>a</sup>	1.21±0.76 <sup>a</sup>	1.93±0.43 <sup>a</sup>
Group 1	2.99±0.35 <sup>b</sup>	0.89±0.11 <sup>a,b</sup>	1.47±0.41 <sup>a,b</sup>
Group 2	2.15±0.32 <sup>b</sup>	0.92±0.52 <sup>a,b</sup>	1.39±0.43 <sup>a,b</sup>
Group 3	3.02±0.07 <sup>a</sup>	1.18±0.16 <sup>a</sup>	1.31±0.54 <sup>a</sup>

Values are expressed as mean ± SD.- Significant at p<0.05 using one way ANOVA test.- Values which have different letters in each column differ significantly, while those with have similar or letters are not significant

**Serum lead, iron and blood hemoglobin of rats suffering from lead toxicity.**

The concentration of lead and iron in serum and blood hemoglobin were determined and the results are reported in Table (5). The concentration of lead in serum was increased significantly when rats were exposed to lead toxicity 41.13 mg/dl (control positive) compared with negative control group was 29.29 mg/dl. When rats fed 15% mixture of banana and mango peels significant reduction in serum lead level compared with positive control group which was close to the normal rats control group. These results were agreement with **Esfandiar and Mahmoud (2012) and El-Nahal(2010)** they showed that, serum lead levels decreased significantly (p< 0.05) in the artichoke- treated group compared to lead-intoxicated rats without treatment.

While the serum concentration of iron and the value of hemoglobin in blood were significantly decreased in control positive (26.11 and 9.79 mg/dl) than the negative control group was 38.93 and 12.91 mg/dl. The rats fed on mixture of banana and mango peels was significant increased in serum iron level and hemoglobin in blood (33.91 and 12.96 mg/dl) than positive control group. Which are similar results to that recorded by **Khanet al.(2008) and El-Nahal(2010)** who showed that the lead acetate administration lead to reduced Hb and PCV. Similarly, **Szymbek et**

al.(1983) observed that Hb level was reduced after intoxication with lead acetate in dose of 400 mg / kg of the fodder. The results were agreed with that obtained by **Ali and Blunden (2003)** who reported that treatment of rats with the black cumin seeds extract for 12 weeks has induced an increased in the hemoglobin level.

**Table (5): Serum lead, iron and blood hemoglobin of rats suffering from lead toxicity.**

Groups	Serum lead (mg/dl)	Serum iron(mg/dl)	Blood hemoglobin (mg/dl)
Control negative	29.29±5.07 <sup>b</sup>	38.93±4.71 <sup>a</sup>	12.91±1.50 <sup>a</sup>
Control positive	41.13±5.74 <sup>a</sup>	26.11±5.38 <sup>b</sup>	9.79±0.40 <sup>b</sup>
Group 1	31.42±3.45 <sup>a,b</sup>	32.87±7.06 <sup>b</sup>	11.94±1.05 <sup>a,b</sup>
Group 2	32.81±2.41 <sup>a,b</sup>	31.73±1.06 <sup>b</sup>	11.54±0.68 <sup>a,b</sup>
Group 3	29.52±0.32 <sup>b</sup>	33.91±5.08 <sup>a</sup>	12.96±1.06 <sup>a</sup>

Values are expressed as mean ± SD.- Significant at  $p < 0.05$  using one way ANOVA test.- Values which have different letters in each column differ significantly, while those with have similar or letters are not significant

**Calcium, phosphorus and lead in Tibia of rats suffering from lead toxicity**

Table (6) revealed that the results concentration of both calcium and phosphorus in tibia rats experimental were significantly decreased when rats were exposed to lead toxicity (control positive) 6.11, 5.11g/100g and 443mg/kg compared with negative control group were 9.81, 8.16 g/100g and 0.311mg/kg, respectively. The rats fed on mixture from banana and mango peel the results showed that significant increased in calcium and phosphorus concentration in tibia 8.50 and 7.16 g/100g compared with the positive control group. On the contrary lead concentration in bone was significant decreased (0.351 mg/kg) compared with rats fed on banana and mango peels was 0.361 and 0.390 mg/kg, respectively. These results were agreement with **Abd -Elhalim et al. (2008)** , **Hamilton and Flaherty (1994)** and **Bagchi and Preuss (2005)** who found that lead accumulation in rats has caused reduction in femoral bone mass density (BMD) compared with the untreated rats with lead intoxicated of similar age. **Imran et**

**al.(2011),Chen et al. (2010) and Naser et al. (2015)** concluded that banana and mango peels improve the concentration of calcium, phosphorous in bone as well as lead in bone of intoxicated rats

**Table (6): Calcium, phosphorus and lead in Tibia of rats suffering from lead toxicity.**

Groups	Calcium (g/100g)	Phosphorus (g/100g)	Lead (mg/Kg)
Control negative	9.81±1.00 <sup>a</sup>	8.16±0.96 <sup>a</sup>	0.311±0.465 <sup>b</sup>
Control positive	6.11±0.30 <sup>b</sup>	5.11±2.05 <sup>b</sup>	0.443±0.510 <sup>a</sup>
Group 1	7.80±0.50 <sup>a,b</sup>	6.18±1.80 <sup>a</sup>	0.361±0.115 <sup>a,b</sup>
Group 2	7.11±0.10 <sup>a,b</sup>	6.00±1.10 <sup>a</sup>	0.390±0.345 <sup>a,b</sup>
Group 3	8.50±1.00 <sup>a</sup>	7.16±0.70 <sup>a</sup>	0.351±0.362 <sup>b</sup>

Values are expressed as mean ± SD.- Significant at p<0.05 using one way ANOVA test.- Values which have different letters in each column differ significantly, while those with have similar or letters are not significant

**Effect of banana and mango peels on serum liver enzymes of rats suffering from lead toxicity.**

Results in Table (7) showed that rats were exposed to lead toxicity control positive was significantly increased level of aspartate amino transferase(AST) and alanine amino transferase (ALT) in serum (104.92 and 35.94 U/L) compared with negative control group was 65.00 and 12.92 U/L, respectively. The resultant in the same table revealed that the rats fed on banana peels;mangopeels and mixture of them were reduction in the elevated serum activity of AST and ALT level compared with positive group. These results agreement with ( **Khan et al., 2008, Saeed .,2015 and El - Tantawy ., 2015**) who reported that, the activities of ALT and AST in serum were significantly increased in lead exposed rats(**Mosa and Khalil, 2015, El Makawy et al , 2015**)showed that banana peelsand mango peels have good protective effects against acute hepatic injury in rats.

**Table (7):- Effect of banana and mango peels on serum liver enzyme of rats suffering from lead toxicity.**

Groups	Serum AST(U/L)	Serum ALT(U/L)
Control negative	65.00±11.10 <sup>c</sup>	12.92±3.63 <sup>c</sup>
Control positive	104.92±3.11 <sup>a</sup>	35.94±26.79 <sup>a</sup>
Group 1	85.67±24.51 <sup>b</sup>	21.10±17.54 <sup>b</sup>
Group 2	89.61±31.14 <sup>b</sup>	25.34±17.11 <sup>a,b</sup>
Group 3	82.11±51.10 <sup>b</sup>	19.61±13.17 <sup>b</sup>

Values are expressed as mean ± SD.- Significant at  $p < 0.05$  using one way ANOVA test.- Values which have different letters in each column differ significantly, while those with have similar or letters are not significant

**Effect of banana and mango peels on serum kidney functions of rats suffering from lead toxicity.**

Results in Table (8) also showed that increased in the concentration of uric acid, urea and creatinine in positive group (2.87, 147.92 and 0.98 mg/dl) than negative control group was 1.82, 97.21 and 0.57 mg/dl, respectively. Whereas, the rats fed banana peels, mango peels and combination of them were significant reduction and the best result group was fed on mixture banana and mango peels. These results are agreement with **El-Shenawy et al. (2009)**, **Nabil et al. (2013)**, **Abd El-Ghany et al. (2015)** and **Salah et al. (2015)** they reported that there were significant increased in serum creatinine, urea, uric acid and globulin of lead intoxicated rats. The results from this study revealed that rats treated with the aqueous extracts of the three cultivars of *Musa acuminata* pose no threat to the liver. Conversely, aqueous extracts of *Saro* cultivars showed increase in the biomarkers of the kidney. Thus, intake of peel extracts of *M. acuminata*, particularly the *Saro* cultivar, as a drug might lead to potential kidney problems in the management and/or treatment of hypertension and other cardiovascular related diseases (**ChidiEdenta et al, 2017**).

**Table (8):- Effect of banana and mango peels on serum kidney functions of rats suffering from lead toxicity.**


Groups	Uric acid (mg/dl)	Urea (mg/dl)	Creatinine (mg/dl)
Control negative	1.82±0.26 <sup>c</sup>	97.21±16.10 <sup>d</sup>	0.57±0.27 <sup>c</sup>
Control positive	2.87±0.71 <sup>a</sup>	147.92±43.60 <sup>a</sup>	0.98±0.47 <sup>a</sup>
Group 1	2.00 ±0.04 <sup>a,b</sup>	104.91±4.00 <sup>c</sup>	0.64±0.07 <sup>c,b</sup>
Group 2	2.01±0.034 <sup>a</sup>	109.05±83.10 <sup>c</sup>	0.68±0.08 <sup>c</sup>
Group 3	1.81±0.27 <sup>c</sup>	103.12±3.11 <sup>c</sup>	0.63±0.53 <sup>b, c</sup>

Values are expressed as mean ± SD.- Significant at p<0.05 using one way ANOVA test.- Values which have different letters in each column differ significantly, while those with have similar or letters are not significant


**Histopathological results:**

The histopathological results were agreed and in same line with the results of serum parameters.


**Histopathological Changes of liver:**




**Photo. (1):**Liver of control -ve rat showing the normal histological structure of hepatic lobule (Hand E X 200 ).




**Photo. (2):**Liver of control +ve rat showing focal area of hepatic necrosis completely replaced by leucocytic cells infiltration (Hand E X 200).



**Photo. (3):** Liver of intoxicated rat fed on diet containing "15% banana peels" showing no histopathological changes (Hand E X 200).




**Photo(4):** Liver of intoxicated rat fed on diet containing "5% mango peels" showing dilatation and congestion of central vein (Hand E X 200).




**Photo (5):** Liver of intoxicated rat fed on diet containing "7.5% banana peels and 7.5 mango peels" showing no histopathological changes (Hand E X 200).


### **Histopathological Changes of bone:**




**Photo. (6):** Bone of rat from control negative group fed on basal diet showing no histopathological changes (H & E X 200).




**Photo. (7):** Bone of rat exposed to lead toxicity as "positive control group" showing thin trabecular bones and enlarged medullary cavity (H & E X 200).



**Photo. (8):** Bone of intoxicated rat fed on diet containing 15% banana peels showing normal thickness cortical bone (H & E X 200).



**Photo. (9):** Bone of intoxicated rat fed on diet containing 15% mango peels showing no histopathological changes (H & E X 200).



**Photo. (10):** Bone of intoxicated rat fed on diet containing 15% mix. of all dried peels showing no histopathological changes (H & E X 200).

**Conclusion and recommendations:**

The present study is scientifically proved that dried banana peels and dried mango peels intake produced great therapeutic effects against lead toxicity and improvements serum lead, iron, blood hemoglobin, calcium, phosphorous, bone, liver enzyme activities and kidney function. The best results showed in group fed diet contain (15%) mix of dried banana and mango peels compare another groups .They could be considered a cheapest nutritious food supplement for sensitive groups .Raising the nutrition awareness with such fruit type for its qualities and biologically effects, it deserves and paying attention for its cultivation, marketing, storage and technological food industries.



## **References**

- Abd El- Ghany, M., sedky, A and El- Metwally, D. (2015):** Effect of consumption different legumes in diets fortified with black cumin and guava seeds on Nephrotoxicity in rats. *Current Science International*.4:45- 55.
- Abd El-Halim, A.,Moustafa, T, Mohamed ,M.,and Ahmed,A.(2008):** Effect of soybean on bone and gonad hormones in lead intoxicated rats. *journal of Biological Research- Thessaloniki*.9:25-34.
- Adejuyitan, J.A., Adelakun, O.E., Olaniyan, S.A., Popoola, F.I., (2008):**Evaluating the quality characteristics of kunun produced from drymilled sorghum. *Afr. J. Biotechnol.* 7 (13), 2244–2247.
- Adlin, N.M.D.Y., 2008.** Correlation between total phenolics and mineral content with antioxidant activity and determination ofbioactive compounds in various local bananas (*Musa sp.*). Thesis, UniversitasDiponegoro, Semarang
- Ajila C.M., Aalami M., Leelavathi K., PrasadaRao U.J.S.,(2010):** Mango peel powder: A potential source of antioxidant and dietary fiber in macaroni preparations. *Inn.Food Sci. Emerg. Technol.*, 2010, 11, 219–224.
- Ajila C.M., PrasadaRao U.J.S.,(2008)** Protection against hydrogen peroxide induced oxidative damage in rat erythrocytes by *Mangifera indica*L. peel extract. *Food Chem. Toxicol.*, 2008, 46, 303–309.
- Ajila C.M., Bhat S.G., PrasadaRao U.J.S., (2007)** Valuable components of raw and ripe peels from two Indian mango varieties. *Food Chem.*, 2007, 102, 1006–1011.
- Ali,B. and Blunden,G.(2003):** Pharmacological and Toxicological Properties of *Nigella sativa*. *Phytotherapy Research.* 17: 299 - 305.
- Anhwange, B.A., Ugye, T.J., Nyiaatagher, T.D., (2008):** Chemical composition of *Musa sapientum* (banana) peels. *Electron. J. Environ. Agric. Food Chem.* 8 (6), 437–442.
- A.O.A.C., 2005.** In: *Official Methods of Analysis*, vol. 8. Association of Official Analytical Chemists, Arlington, Virginia, D.C., USA,pp. 242–273.

- AOAC (2010).** Official Methods of Analysis of Association of Official Chemists. 18<sup>th</sup> Ed., Washington, D.C., USA.
- Ashoush, I.S., & Gadallah, M.G.E . (2011).** Utilization of Mango Peels and Seed Kernels Powders as Sources of Phytochemicals in Biscuit. *World Journal of Dairy and Food Sciences*, 6(1), 35-42.
- Bagchi D, Preuss H. (2005):** Effect of acute and chronic oval exposure of lead on blood pressure and bone mineral density in rats. *Journal of inorganic biochemistry*, 99: 1155-1164.
- Carter M. and Gregorich.E.(2006):** Soil Sampling and Methods of analysis 2nd ed:P. cm. vol. 13: 3586 - 8493.
- Cao, G., Booth, S. L., Sadowski, J. A., & Prior, R. L. (1998).** Increases in human plasma antioxidant capacity after consumption of controlled diets high in fruit and vegetables. *The American journal of clinical nutrition*, 68(5), 1081-1087.
- Carbonneau, M. A., Léger, C. L., Descomps, B., Michel, F., & Monnier, L. (1998).** Improvement in the antioxidant status of plasma and low-density lipoprotein in subjects receiving a red wine phenolics mixture. *Journal of the American Oil Chemists' Society*, 75(2), 235-240.
- Chapman, D.G.; Gastilla, R. and Campbell, J.A. (1959):** Evaluation of protein in food. I. A. Method for the determination of protein efficiency ratio. *Can J Biochem Physiol.*, 37:679-686.
- Chen, J.R.; Lazarenko, O.P.; Wu, X.; Kang, J.; Blackburn, M.L.; Shankar, K.;**
- Badger, T.M. and Ronis, M.J. (2010):** Dietary-induced serum phenolic acids promote bone growth via p38 MAPK/ $\beta$ -catenin canonical Wnt signaling. *J Bone Miner Res.*, 25(11):2399-411.
- Chidi Edenta, Stanley I. R. Okoduwa, ID and Oche Okpe(2017):** Effects of Aqueous Extract of Three Cultivars of Banana (*Musa acuminata*) Fruit Peel on Kidney and Liver Function Indices in Wistar Rats. [www.mdpi.com/journal/medicines](http://www.mdpi.com/journal/medicines) doi: 10.3390.
- Davis, J. and Svendsgaard, D. (1987):** lead and child development. US Environmental Protection Agency, Research Triangle Park, North Carolina. 329(7): 297 - 300.

- El-Nahal, M. D. (2010):** Effect of using pectin on lead toxicity The Journal of American Science,2010; 6(12) <http://www.americanscience.org>.
- El- Shenawy, N., Al- Eisa, R., El- Salmy, F. and Salah,O.(2009):** Prophylactic effect of vitamin E against hepatotoxicity, nephrotoxicity, haematological induces and histopathology induced by diazinon insecticide in mice. *Curr.Zool.*55(3):219- 226.
- El- Tantawy, W, H (2015):** Antioxidant effects of Spirulina supplement against lead acetate - induced hepatic injury in rats. Original article.doi: 10. 1016/j. jtcme.02.001.
- El Makawy Aida I, Ihab S. Ashoush, Omaima M. Abd-Elmoneim(2015):** Evaluation of Mango Byproduct Extracts as Antioxidant Against Pb-Acetate-Induced Oxidative Stress and Genotoxicity in Mice . *J. Food Nutr. Sci.*, 2015, Vol. 65, No. 1, pp. 39–47.
- Ercal, N., Treeratphen, P., Hammond, T., Mattews, R., Grnnemann, N. and Spitz, D. (1996):**In vivo indices of oxidative stress in lead exposed c57bl/6 mice reduced by treatment with meso 2, 3-dimercaptosuccinic acid or nacetylcystein.*FreeRadic. Biol.Med.* 21: 157- 161.
- Esfanddiar,H, Mahmoud .R, K.(2012):** Protective effect of artichoke (*Cynarascolymus*) leaf extract against lead toxicity in rats. 51: 1104 - 1109.
- FAO,(2012):** Food and Agriculture Organization, FAOSTAT database collections, agricultural data, food and agriculture organization of the United Nations. Available from: [<http://faostat.fao.org>], 2012.
- Faigin, R., (2001):**MeningkatkanHormonSecaraAlami. PT Raja GrafindoPersada, Jakarta.Figuerola, F., Hurtado, M.L., Estevez, A.M., Chiffelle, I., Asenjo, F., 2005.Fibre concentrates from apple pomace and citrus peel as potential fibre sources for food enrichment. *Food Chem.* 9, 395–401
- Figuerola, F., Hurtado, M.L., Estevez, A.M.,Chiffelle, I., Asenjo, F., (2005):**Fibre concentrates from apple pomace and citrus peel as potential fibre sources for food enrichment. *Food Chem.* 9, 395–401.

- Garcia-Salas P, Morales-Soto A, Segura-Carretero A, Fernández-Gutiérrez A. 2010.** Phenolic compound extraction systems for fruit and vegetable samples. *Molecules*, 15:8813-8826
- Gonzalez-Montelongo, R.M., Gloria Lobo, G., Gonzalez, M., 2010.** Antioxidant activity in banana peel extracts: testing extraction conditions and related bioactive compounds. *Food Chem.* 119,1030–1039.
- HappiEmaga, T., Andrianaivo, R.H., Wathelet, B., Tchango, J.T., Paquot, M., 2007.** Effects of the stage of maturation and varieties on the chemical composition of banana and plantain peels. *Food Chem.* 103, 590–600.
- Haisman, P. and Muller, B. R. (1977):** Quantitative enzymatic colorimetric determination of uric acid in serum. *Clin Chem.*, 26: 227.
- Hamilton, J. and Flaherty, E. (1994):** Effects of lead exposure on skeletal development in rats. *Fundamental and applied toxicology*, 22:594-604.
- Hertog, M. G. L., Kromhout, D., Aravanis, C., Blackburn, H., Buzina, R., Fidanza, F., Ciampaoli, S., Jansen, A., Menotti, A., Nedeljkovic, S., Pekkarinen, M., Simic, B. S., Toshima, H., Feskens, E. J. M., Hollman, P. C. H., & Katan, M. B. (1995).** Flavonoid intake and long-term risk of coronary heart disease and cancer in the seven countries study. *Arch. Int. Med.* 155(4), 381-386.
- Hertog, M. G., Feskens, E. J., Kromhout, D., Hollman, P. C. H., & Katan, M. B. (1993).** Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen Elderly Study. *The Lancet*, 342(8878), 1007-1011.
- Henry, R. (1974):** Determination of creatinine by kinetic method. In "Clinical Chemistry, Principles and techniques "( 2<sup>nd</sup> edition), Harper & Rowe, 525.
- Howarth, N.C., E. Saltzman and S.B. Roberts, 2001.** Dietary fiber and weight regulation. *Nutr. Rev.*, 59(5): 129-139.

- Imran, A.; Jat, R.K. and Varnika, S. (2011):** A Review On Traditional, Pharmacological, Pharmacognostic Properties Of FicuCarica (Anjir). *IRJP.*, 2 (12): 124- 127.
- Ismail, F., Anjum, M. R., Mamon, A. N. and Kazi, T. G. (2011)***Trace metal contents of vegetables and fruits of Hyderabad retail market*, Pakistan Journal of Nutrition, 10, 365–372.
- Kanazawa, K., Sakakibara, H., (2000):** High content of dopamine, a strong antioxidant, in Cavendish banana. *J. Agric. Food Chem.* 48 (3), 844–848.
- Karamian, R., Komaki, A., Salehi, I., Tahmasebi, L., Komaki, H., Shahidi, S., & Sarihi, A. (2015).** Vitamin C reverses lead-induced deficits in hippocampal synaptic plasticity in rats. *Brain research bulletin*.
- Khan, M., Mostofa, M., Jahan, M., Sayed, M. and Hossain, M. (2008):** Effect of garlic and vitamin B-complex in lead acetate induced toxicities in mice. *Bangl. J. V. Med.*, 6(2): 203-210.
- Knekt, P., Jarvinen, R., Reunanen, A., & Maatela, J. (1996).** Flavonoid intake and coronary mortality in Finland: a cohort study. *Bmj*, 312(7029), 478-481.
- Khodamoradi, N., Komaki, A., Salehi, I., Shahidi, S., & Sarihi, A. (2015).** Effect of vitamin E on lead exposure-induced learning and memory impairment in rats. *Physiology & behavior*, 144, 90-94.
- Kondo, S., Kittikorn, M., Kanlayanarat, S., (2005):** Preharvest antioxidant activities of tropical fruit and the effect of low temperature storage on antioxidants and jasmonates. *Postharvest Biol. Technol.* 36, 309–318.
- Larson, R. A. (1988).** The antioxidants of higher plants. *Phytochemistry*, 27, 969-978.
- Larsen, K. (1972).** Creatinine colorimetric method. *J. of Clin. Chem.*, 41:209.
- Li, X., Ye, F., Li, L., Chang, W., Wu, X., and Chen, J. (2016).** The role of hemo-1 in protection against lead-induced neurotoxicity. *Neurotoxicology*, 52, 1-11.


- Llobera, A., Canellas, J., (2007):** Dietary fibre content and antioxidant activity of Manto Negro red grape (*Vitisvinifera*): pomace and stem. *Food Chem.* 101, 659–666.
- Manthey J.A., Perkins-Veazie P.(2009):**, Influences of harvest date and location on the levels of  $\beta$ -carotene, ascorbic acid, total phenols, the *in vitro* antioxidant capacity, and phenolic profiles of five commercial varieties of mango (*Mangifera indica* L.). *J. Agri. Food Chem.*, 2009, 57, 10825–10830.
- Marin, F.R., Soler-Rivas, C., Benavente-Garcia, O., Castillo, J., Perez-Alvarez, J.A., (2007):** By-products from different citrus processes as a source of customized functional fibres. *Food Chem.* 100, 736–741.
- Matook, S.M.I., Fumio, H. (2005):** Antibacterial and antioxidant activities of banana (*Musa*, AAA cv. Cavendish) fruits peel. *Am. J. Biochem. Biotechnol.* 1 (3), 125–131.
- Marchlewicz, m.; m. Protasouicki; I. Rozewicka; m. Piasecka and m. Laszczynska (1993).** Effect of long-term exposure to lead on testis and epididymis in rats. *Folia-histochem.Cytobiol.* 31 (2):55–62.
- Mingeum, J., Dong, S. L., Hyung, J. S. and Yooheon, P. (2015):** *Aspergillus Oryzae* fermented germinated soybean extract alleviates perimenopausal symptoms in ovariectomised rats. Article first published online: 23.
- Mosa Z.M., Khalil A. F (2015):** The effect of banana peels supplemented diet on acute liver failure rats. *Annals of Agricultural Science* 60(2), 373–379.
- Nabil, T., Mahdy, K., Abdel- Wahab, M. and Mohamed, L.(2013):** Effect of lead toxicity on mineral metabolism and immunological factors in rats. *Alexandria Journal of Veterinary Sciences.* 39(8):64 - 73.
- Naser. B. A ,El-Bahy. A. M, Abd El-Aziz. A, Mohamed H. N and Ibrahim M. Aniess.W.( 2015):** Effect of Soybean and Some Dried Fruits in Osteoporotic Rats. M.Sc. In Education Faculty of Education - Ain Shams University.

- Newairy, A. and Abdou, H. (2009):** Protective role of flax lignans against lead acetate induced oxidative damage and hyperlipidemia in rats. *Food Chem Toxicol.* 47( 4): 813-818.
- Parsons ,P (2001):** Analytical procedures for the determination of lead in blood and urine, approved guideline. Wayne, PA, National Committee for Clinical Laboratory Standards. 113 (8):1669- 1647.
- Patrick, L. (2006).** Lead toxicity part II: the role of free radical damage and the use of antioxidants in the pathology and treatment of lead toxicity. *Altern. Med. Rev.*11: 114–127.
- Ramsay W.N.M. 1957.** The determination of iron in blood plasma and serum *Clin. Chem. Acta.*2:214-221.
- Reeves, P.G.; Nielsen, F.H. and Fahey, G.C. (1993):** AIN-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the AIN-76A rodent diet. *J Nutr.*, 123(11):1939-51.
- Rice-evans, C. A., Miller, N. J., Bolwell, P. G., Bramley, P. M., & Pridham, J. B. (1995).** The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free radical research*, 22(4), 375-383.
- Saeed A. A. (2015):**Haemato - biochemical changes induced by lead intoxication in male and female albino mice. *International Journal of Recent Scientific Research.*6(5):2999 - 4004.
- Salah, A., Zadjali, N, Mohamed,Y., Fahim, S, Azimullah ,A,Subramanian,D,Yasin, Y., Amir,N,Hassan,M, and Adem, A. (2015):**Lead exposure causes thyroid abnormalities in diabetic rats. *Int J Clin EXP Med.*8(5): 7160- 7167.
- SAS System for Windows (Statistical Analysis System) (2008).**Version 9.2. Cary, USA: SAS Institute Inc. S.
- Schieber A., Berardini N., Carle R.,(2003):**Identification of flavonol and xanthone glycosides from mango (*Mangifera indica*L. Cv. ‘‘Tomy Atkins’’) peels by high performance liquid chromatography–electrospray ionizations mass spectrometry. *J. Agr. Food Chem.*, 51, 5006–5011.

- Serafini, M., Maiani, G., & Ferro-Luzzi, A. (1998).**Alcohol-free red wine enhances plasma antioxidant capacity in humans. *The Journal of Nutrition*,128(6), 1003-1007.
- Silbergeld, E. K; M. Waalkes and Rice, J. M. (2000).** Lead as a carcinogen: experimental evidence and mechanisms of action. *Am. J. Ind. Med.* 38: 316–323.
- Sherwin, J.(1984):** Liver function. In: Kaplan LA, pesce AJ eds, clinical chemistry, theory, analyses, and correlation. Vol. 3: 420- 438.
- Sogi, D. S., Siddiq, M., Greiby, I., & Dolan, K. D. (2013).**Total phenolics, antioxidant activity, and functional properties of ‘Tommy Atkins’ mango peel and kernel as affected by drying methods. *Food chemistry*, 141(3), 2649-2655.
- Someya, S., Yoshiki, Y., Okubo, K., (2002):** Antioxidant compounds from bananas (*Musa Cavendish*). *Food Chem.* 79, 351–354.
- Subagio, A., Morita, N., Sawada, S.(1996):** Carotenoids and their fatty-acid esters in banana peel. *J. Nutr. Sci. Vitaminol.* 42 (6), 553–566.
- Sulaiman, S.F., Yusoff, N.A.M., Eldeen, I.M., Seow, E.M., Sajak, A.A.S., Ooi, K.L., (2011):** Correlation between total phenolic and mineral contents with antioxidant activity of eight Malaysian bananas (*Musa sp.*). *J. Food Compos. Anal.* 24, 1–10.
- Szymezak,J., Zechalko, A. and Biernat,J (1983):** Effect of foodder fat type on blood plasma lipids in mice intoxicated with lead. *BromatolChemToksykol* 16(2): 98 -94.
- Tetens, I. and S. Alinia, (2009):** The role of fruit consumption in the prevention of obesity. *Journal of Horticultural Science & Biotechnology (Special Issue):* 47-51.
- The World Health Report (2015):** The double burden: Emerging epidemics and persistent problems. WHO 2015. Available at: <<http://www.who.org>>.
- Willett, W.C., F. Sacks and A. Trichopoulou, 1995.** Mediterranean diet pyramid: a cultural model for healthy eating. *Am. J. Clin. Nutr.*, 61: 1402S-1406S.



- World Health Organization (WHO) statistics the Institute for Health Metrics and Evaluation ( IHME) 2013:** GBD compare, Seattle, WA: IHME, University of Washington.
- Xu BJ. And Chang SKC. (2007):** A comparative study on phenolic profiles and antioxidant activities of legumes as affected by extraction solvents. *J Food Sci*, 72:159-166.
- Young, D. (1990):** Effects of drugs on clinical laboratory test. Third Edition. 3:6-12.



## التاثير الوقائى لقشور الموز و المانجو الجافه ضد سميته الرصاص فى الجرذان

ايرينى ولسن نجيب , هاله راشد عطايا

قسم الاقتصاد المنزلى - كلية التربية النوعية- جامعة عين شمس

### الملخص العربى

تهدف هذه الدراسة لمعرفة تأثير اضافته كلا من قشر الموز و قشر المانجو المجففان للوجبه الأساسية ضد سميته الرصاص فى الجرذان. تم اجراء الدراسة على عدد (45) جرذ من نوع ذكر الالبينو البياض فى فترة النمو وتم تقسيمهم الى مجموعتين رئيسيتين. المجموعة الاولى (المجموعة الضابطة السالبة) (٩) جرذان. اما المجموعة الثانية (36) جرذ قد تعرضت للتسمم بالرصاص و قسمت الى اربع مجموعات بكلا منها (٩) جرذان. المجموعة الاولى تغذت على الغذاء الاساسى (المجموعة الضابطة الموجبه) و الثانية تغذت على الغذاء الاساسى يحتوى على (١٥%) قشر موز , الثالثة تغذت على الغذاء الاساسى يحتوى على (١٥%) قشر مانجو , الرابعة تغذت على الغذاء الاساسى يحتوى على (١٥%) خليط (٧.٥% قشر موز + ٧.٥% قشر مانجو) لمدة (٦) اسابيع. اثناء فترة التجربة تم حساب كلا من معدل الزيادة فى وزن الجسم , كمية الطعام المتناول و كذلك معدل كفاءة الغذاء و فى نهاية التجربة تم تجميع عينات الدم وتقدير هيموجلوبين الدم ثم تقدير كلا من الحديد , الرصاص , انزيمات الكبد , ووظائف الكلى فى السيرم و تقدير كلا من مستوى الكالسيوم و الفسفور و الرصاص فى العظام. اظهرت النتائج ان الجرذان التى تناولت خليط قشور الموز و المانجو كانت تمثل افضل النتائج فى معدل الزيادة فى الوزن و الطعام المتناول و معدل كفاءة الطعام و تحسين فى مستوى الهيموجلوبين و الحديد و انزيمات الكبد ووظائف الكلى و حدوث انخفاض فى مستوى الرصاص فى السيرم مقارنة بالمجموعة الضابطة الموجبة و اكدت نتائج الفحص الهستوباثولوجى نفس النتائج السابقة. لذلك توصى الدراسة بعمل برامج للتوعية الغذائية لتوضيح اهمية قشور الموز و المانجو فى خفض خطر الاصابه أو التلوث بالتسمم بالرصاص.

**الكلمات المفتاحية:** التسمم بالرصاص , قشر الموز , قشر المانجو.